

Feelnor®

Trimetazidine Dihydrochloride Tablet

Presentation

Feelnor 20 tablet: Each film-coated tablet contains Trimetazidine Dihydrochloride BP 20 mg.

Feelnor 35 MR tablet: Each modified release tablet contains Trimetazidine Dihydrochloride BP 35 mg.

Description

Trimetazidine is a unique antianginal and anti-ischæmic agent that display anti-ischæmic effects without inducing haemodynamic changes and improves the status of the ischæmic myocardium. The drug exerts its antianginal effects through a direct cytoprotective action on the myocardium, thereby avoiding the side-effects of impaired left ventricular function, excessive peripheral vasodilation and other disadvantages associated with the use of other antianginal agents.

Indications and Uses

Trimetazidine 20 mg is indicated for-

- Ischæmic heart disease (angina pectoris, sequelae of infarction)
- Adjuvant symptomatic treatment of vertigo and tinnitus
- Adjuvant treatment of the decline visual acuity and visual field disturbances, presumably of vascular origin

Trimetazidine 35 mg MR is indicated for-

- Long-term treatment of angina pectoris

Dosage & Administration

One 20 mg tablet thrice daily after meals. No dosage adjustments are required in patients with impaired renal and hepatic function.

One 35 mg MR tablet twice daily at mealtimes in the morning and evening.

Side-effects

Trimetazidine is safe and well tolerated. The most commonly side effects are gastric discomfort, nausea, headache and vertigo. However, the side effects are mild and non-specific.

Use in Pregnancy & Lactation

Pregnancy: Studies in animals have not demonstrated a teratogenic effect. However, in the absence of clinical data and for safety reasons, prescription should be avoided during pregnancy.

Nursing Mothers: There is no information on the secretion of Trimetazidine into breast milk. However, breast feeding should be discontinued if the use of Trimetazidine is considered essential.

Pediatric Use

Safety and efficacy in children have not been established with Trimetazidine.

Contraindications

Hypersensitivity to Trimetazidine Dihydrochloride.

Drug Interactions

No drug interactions have so far been reported. In particular, no interactions of Trimetazidine with beta-blockers, calcium antagonists, nitrates, heparin, hypolipidemic agents or digitalis have been reported.

Commercial Pack

Feelnor 20: Each box contains 3 blister strips of 10 tablets.

Feelnor 35 MR: Each box contains 3 blister strips of 10 tablets.

Manufactured by
 **Incepta Pharmaceuticals Ltd**
Dhaka, Bangladesh
® Registered Trademark

FEM