



Myolax[®]

Tolperisone Hydrochloride

Presentation

Myolax[®] 50: Each film coated tablet contains Tolperisone Hydrochloride INN 50 mg.

Description

Tolperisone Hydrochloride is a centrally acting muscle relaxant. It effectively reduces muscular hypertonicity, improves peripheral blood flow by vasodilatation and reduces pain.

Indications and uses

Increased tone of skeletal muscles due to organic neurological disorders (e.g. injury of the pyramidal tract, multiple sclerosis, myelopathy, encephalomyelitis etc.), muscular hypertension, muscular spasm, muscular contracture, rigidity, spinal automatism and discopathy. Obliterative vascular diseases (obliterative arteriosclerosis, diabetic angiopathy, obliterative thromboangitis, Raynaud's disease, diffuse scleroderma) and disorders due to injured innervation of the vessels (acrocyanosis, intermittent angioneurotic dysbasia). In individual cases post-thrombotic venous and lymphatic circulation disorders, crural ulcer.

Dosage and Administration

One should follow doctor's instructions regarding the dosage of Myolax. The recommended dose is as follows:

Tablet

Adult: 50-150 mg three times daily according to the individual requirement and tolerance of the patient.

Children (from 3 months to 6 years): 5-10 mg/kg/day in three divided doses.

Children (from 6 to 14 years): 2-4 mg/kg/day in three divided doses.

Contraindications

- Hypersensitivity to any component of the product
- Myasthenia gravis
- Nursing mother

Precautions

In case of children, the prescribed dose and duration of treatment should closely be observed.

Side-effects

Sometimes, muscular weakness, headache, nausea, vomiting and rarely hypersensitivity reactions may occur as side effects.

Use in Pregnancy and Lactation

Tolperisone hydrochloride should be used in pregnancy according to physician's advice. Tolperisone hydrochloride should not be used during breast feeding.

Drug Interactions

Concomitant use of methocarbamol with tolperisone hydrochloride has been reported to cause disturbance of visual accommodation.

Over dosage

In the event of over dosage supportive therapy may be substituted.

Storage

Do not store above 30°C. Keep away from light and out of the reach of children.

Commercial Pack

Myolax[®] 50: Each box contains 7 blister strips of 10 tablets.

Manufactured by
 **Incepta Pharmaceuticals Ltd**
Dhamrai Unit, Dhaka, Bangladesh
® Registered Trademark

VN.01

MY2