

Netario™ Sterile Eye Drops

Netarsudil 0.02%

Presentation

Netario™ Sterile Eye Drops: Each ml eye drops contain Netarsudil dimesylate INN equivalent to Netarsudil 0.2 mg.

Description

Netarsudil is both a Rho kinase and norepinephrine transport (NATs) s inhibitor that specifically targets and inhibits rho kinase and NATS found in the conventional trabecular pathway while many of its contemporaries offer therapy that focuses on cell and muscle tissue remodeling.

Indications

Netarsudil 0.02% Sterile Ophthalmic Solution is a Rho kinase inhibitor indicated for the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension.

Dosage and Administration

The recommended dosage is one drop in the affected eye(s) once daily in the evening. If one dose is missed, treatment should continue with the next dose in the evening. Twice a day dosing is not well tolerated and is not recommended. If Netarsudil is to be used concomitantly with other topical ophthalmic drug products to lower IOP, administer each drug product at least 5 minutes apart.

Side Effects

The most common adverse reaction is conjunctival hyperemia (53%). Other common adverse reactions, approximately 20% include: corneal verticillata, instillation site pain, and conjunctival hemorrhage.

Contraindications

Contraindicated in patients with previously demonstrated hypersensitivity to any of the ingredients in the formulation.

Precaution

There have been reports of bacterial keratitis associated with the use of multiple-dose containers of topical ophthalmic products. Contact lenses should be removed prior to instillation of Netarsudil and may be reinserted 15 minutes following its administration.

Use in Special Group

Pregnancy: There are no available data on Netarsudil use in pregnant women to inform any drug associated risk; however, systemic exposure to netarsudil from ocular administration is low

Lactation: There are no data on the presence of Netarsudil in human milk, the effects on the breastfed infant, or the effects on milk production. However, systemic exposure to netarsudil following topical ocular administration is low.

Pediatric Use: Safety and effectiveness in pediatric patients below the age of 18 years have not been established.

Use in Elderly Patients: No overall differences in safety or effectiveness have been observed between elderly and other adult patients.

Storage

- Store at 2°- 8°C
- Keep out of the reach of children
- Do not use after 6 weeks of opening

Commercial Pack

Netario™ Sterile Eye Drops: Each plastic dropper bottle contains 3ml of sterile eye drops.