Tinabil™

Bilastine

Presentation

Tinabil™ Tablet: Each tablet contains Bilastine INN 20 mg.
Tinabil™ Oral Solution: Each 5 ml oral solution contains Bilastine INN 12.5 mg.

Description

Bilastine is non-sedating, long-acting histamine antagonist. Its effects are mediated via selective inhibition of peripheral H_1-receptors. It shows moderate to high affinity for histamine H_1-receptors and no affinity for muscarinic, serotonergic, dopaminergic and noradrenergic receptors. Bilastine inhibit histamine-induced flare skin reactions for 24 hours following single doses.

Indications and Usage

Bilastine is indicated for the treatment of allergic rhino-conjunctivitis (seasonal and perennial), allergic rhinitis and urticaria.

Dosage and Administration

Adults and adolescents (12 years of age and over): Tinabil™ 20 mg tablet (1 tablet) once daily.
Children (6 to 11 years): Tinabil™ 4 ml oral solution once daily.
The maximum recommended daily dose is 20 mg and should not be exceeded. The tablet should be taken one hour before or two hours after intake of food or fruit juice.

Use in Pregnancy and Lactation

There are no adequate and well-controlled studies in pregnant women. Until such data become available, Bilastine should be avoided during pregnancy, unless advised otherwise by a physician.

Use in Children

The safety and efficacy of Bilastine in children under 12 years of age have not been established.

Side-effects

The most common side effects associated with Bilastine is headache, dizziness and fatigue.

Contraindications

Bilastine is contraindicated in patient who are hypersensitive to this medication or to any of its ingredients.

Drug Interactions

No clinically important drug interactions have been reported.

Overdosage

No serious adverse events have been reported in case of over dosage. However, in the event of overdosage, symptomatic and supportive treatment is recommended.

Storage

Do not store above 30°C. Keep away from light and out of the reach of children.

Commercial Pack

Tinabil™ Tablet: Each box contains 3 Alu-Alu blister strips of 10 tablets.
Tinabil™ Oral Solution: Each bottle contains 60 ml oral solution.

Manufactured by
Incepta Pharmaceuticals Ltd
Dhaka, Bangladesh
™ Trademark

V.N. 01